

Habituation of Predator Inspection and Boldness in the Guppy (*Poecilia reticulata*)*

S. V. Budaev and D. D. Zworykin

Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninskii prospect 33, Moscow, 119071 Russia
e-mail: sbudaev@mail.ru

Received April 15, 2003

Abstract—This study examined habituation of the predator inspection behavior in the guppy (*Poecilia reticulata*) and its relationship with boldness (open field locomotion). Two different strategies were discovered: (1) initial inspection of a predator-like fish, correlated with boldness; (2) subsequent surveillance, governed by a random underlying process and unrelated to boldness. The surveillance inspection is probably linked with anti-predator vigilance. Possible implications to between-population variation in inspection behavior are discussed.

The predator inspection behavior occurs when a fish approaches a potentially dangerous predator-like object. In a typical case, one or several individual fish leave the school, move to the predator, observe it, and then go back (Pitcher *et al.*, 1986; Magurran and Pitcher, 1987).

Approaching a predator involves a high risk of being eaten, and the basic benefits of predator inspection include acquiring information about the predator (Magurran and Girling, 1986; Magurran, 1990), its motivational state (Licht, 1989), and deterring the predator's attacks (Godin and Davis, 1995), although there may be other benefits such as advertising the inspector's quality to potential mates (Godin and Dugatkin, 1997; see Dugatkin and Godin, 1992 for a costs-benefits review). Thus, inspection behavior plays an extremely important role as a predator avoidance mechanism in natural environments (Magurran, 1990). Indeed, fish from populations with a high perdition risk show a higher tendency to inspect predators (Magurran and Seghers, 1990, 1994; Magurran *et al.*, 1992, 1995).

Because predator inspection is dangerous, it may be associated with boldness, i.e., the propensity to take risks (Wilson *et al.*, 1994, Budaev, 1997a). However, the fish from populations with higher predation risk show many signs of shyness¹ (slower habituation of anti-predator responses, more pronounced responses to the alarm substance, higher schooling tendency, etc.) but simultaneously have a higher tendency to inspect (reviewed by Magurran *et al.*, 1992, 1995). Also, sticklebacks (*Gasterosteus aculeatus*) of higher social rank (presumably bolder) showed as low a tendency to inspect as subordinate (presumably the most timid)

ones, and the best inspectors were fish with intermediate rank (McLeod and Huntingford, 1994). Individual differences in predator inspection were not consistent in the guppy (*Poecilia reticulata*) and correlated with exploration but not fear (Budaev, 1997b). Thus, the relationships between predator inspection and boldness are not completely clear: predator inspection might be associated with boldness or shyness. In this study we examined the relationships between predator inspection behavior and boldness in the guppy (*Poecilia reticulata* Peters). We tested how this relationship changed with habituation of the fish to the predator-like object during repeated encounters.

MATERIALS AND METHODS

Twenty-nine adult male guppies (average standard length 1.8 cm) were used in the present experiments. During the experiments, the fish were kept individually in four 100-l aquariums divided by partitions into twelve compartments. The fish were fed daily with commercial fish flakes.

Each individual was tested for boldness in the open field test. In the guppy and several other fish species, open field behavior is correlated with both exploration and lack of fear (Budaev, 1997b; Budaev *et al.*, 1999). This test was performed in a hexagonal tank 0.9 m in diameter, and a 60 W lamp was suspended 1.5 m above. A coordinate grid 10 × 10 cm was marked on the bottom of the tank to record the locomotor activity of the fish. During the testing, the fish was gently released into a white bottomless opaque plastic cylinder, standing in the center of the open field. After two minutes of acclimation, the cylinder was lifted and the behavior of the fish was observed from above for 5 minutes. We recorded locomotor activity of the fish, that is, the number of crosses of the coordinate grid per minute.

¹ Shyness is the opposite of boldness, i.e., the propensity to avoid risk (Wilson *et al.*, 1994).

*This article was submitted by the authors in English.




Fig. 1. Characteristics of the inspection behavior over three days of testing. The percentage of time inspecting and the frequency of inspections are given in logarithmic scale.

One month after the open field test, the guppies were tested in a test of the inspection behavior. It was conducted in an aquarium (60 × 30 × 20 cm) divided into three unequal compartments. The model predator (convict cichlid *Archocentrus nigrofasciatus*), was placed into a small compartment at the flank of the aquarium, separated by a glass wall from the big central compartment. At the opposite end of the aquarium there was a third small compartment, separated by an opaque partition with a door. The tested guppy was released into the last compartment for five minutes, then we opened the door, and when the fish moved to the central compartment, we observed it for five minutes. Each guppy was tested in this test three times during three consecutive days. We recorded (1) the number of predator inspections (frequency per minute), (2) the per-

centage of the time spent inspecting, (3) the percentage of the time spent in proximity (<6 cm) to the compartment with the model predator, and (4) the latency to the first predator inspection visit.

For the statistical analysis, we calculated Friedman ANOVA, sign test, and Spearman correlations (Sokal and Rohlf, 1981). In cases of behavioral latencies, we used statistical methods of survival analysis (Lee, 1992).

RESULTS

The percentage of time spent inspecting and the frequency of predator inspections exponentially reduced with repeated encounters of the cichlid (Figs. 1a, 1b). This is the most commonly observed pattern of decline of a behavioral response to constant stimulus (see Hinde 1970). The time spent in proximity to the model predator also decreased, but only in the last exposure (Fig. 1c): it did not differ significantly from the first to the second exposure, but significantly reduced in the third exposure (the differences with the first two days: sign test, $ps < 0.05$). Thus, the guppies changed their behavior in a way indicating habituation of predator inspection, well documented in the literature (Csanyi, 1985; Magurran and Girling, 1986; Csanyi *et al.*, 1989; Huntingford and Coulter, 1989).

The distribution of the latency to the first inspection significantly deviated from exponential distribution on the first and second days of testing, but not on the third day (Table 1). Kaplan-Meier analysis indicated that, during the first day, the fish tended to inspect the model predator very soon after they entered the central compartment (median latency to inspect = 32 s, i.e., almost 50% of guppies begun to inspect the convict cichlid within the first 1/2 min of the test). The exponential distribution of the inspection latency during the third day means that the probability of the decision to inspect (hazard rate) did not depend on the time elapsed since the guppy entered the compartment with the model predator. The estimates of the hazard rate reduced from the first to the third day of testing (Table 1). In other words, the guppies tended to inspect the convict cichlid rapidly during the first day, but by the third day, the decision to approach the model predator became governed by a single underlying random process, similar to radioactive decay with low probability.

Correlations between the percentage of time inspecting in different days were all in significant ($p > 0.1$, the maximum correlation was between the second and third days, Spearman $r_s = 0.34$, exact $p = 0.11$). Correlations between the frequency of inspections were significant only between the second and third days (Spearman $r_s = 0.40$, exact $p = 0.031$, all other $ps > 0.1$). The percentage of time spent near the compartment with the model predator did not correlate between days (all $ps > 0.1$).

Open field locomotion (boldness) correlated with the inspection behavior measures only in the first day of the inspection testing (Table 2). The same pattern was

also characteristic of the latency to inspect: during the first day of testing, there was a significant positive relationship between the inspection latency and open field locomotion ($\beta = 7.3$, $p = 0.016$, Cox proportional hazard regression), but during the second and third days, this relationship was not significant ($p > 0.1$, Cox proportional regression model).

DISCUSSION

This study showed that predator inspection is a relatively complex behavior in which at least two strategies can be distinguished. The first is exploratory predator inspection, which is associated with boldness and appears first when a predator-like fish is found in proximity. The second strategy of predator inspection is not related to boldness and may be tentatively called surveillance. In the latter case, the fish does not approach and inspect the predator-like fish shortly after it is detected. Rather, the inspection tendency is described by a random process similar to the radioactive decay. Apparently, surveillance looks similar to patrolling (Birke and Archer, 1983; Cowan, 1983) observed in cases of spatial exploration, when an animal is actively searching for changes in a known environment rather than exploring a novel one. In this way, the inspection visits randomly elicited in habituated guppies would be directed to monitoring a possible change of the current state of the potentially dangerous large cichlid, for example, its motivation (Licht, 1989). Also, it may serve the function of signaling that the possible predator has been discovered and still is not forgotten (Hasson, 1991; Godin and Davis, 1995).

The surveillance inspection behavior may suggest that the guppies really perceived the convict cichlid as a potentially dangerous predator. Cichlid predators looking similar are sympatric with many guppy populations in their natural sites in Trinidad (Liley and Seghers, 1975; Reznick and Endler, 1982). This is why this species could be equipped with an inherited mechanism for the recognition of unique features of predators, such as characteristic body shape, which do not disappear even after artificial breeding for several generations (Csanyi, 1985; Altbacker and Csanyi, 1990). Thus, surveillance inspection may reflect antipredator vigilance of the fish to the possible predator (see Lima 1994, 1995 for a discussion on antipredator vigilance). However, it was still not associated with boldness.

Consistency of individual differences in the inspection behavior also reflects two strategies. The initial exploratory inspection, observed during the first day of the experiment and associated with boldness, did not correlate with the surveillance inspection during the subsequent days. However, individual differences in the surveillance inspection tended to be consistent over the last two days. This also suggests that there may be two distinct mechanisms governing approaching of the potential prey to the possible predator. In real situations, there may be complex interactions between the

Table 1. Fitting exponential distribution to the latency of the first inspection

Day of testing	Lambda \pm SE
1	0.0066 ± 0.0022 , $\chi^2 = 23.99$, $df = 8$, $p = 0.002$
2	0.0028 ± 0.0011 , $\chi^2 = 19.90$, $df = 8$, $p = 0.011$
3	0.0008 ± 0.0004 , $\chi^2 = 4.84$, $df = 4$, $p = 0.304$

Note: Lambda (hazard rate) is the parameter of the exponential distribution.

Table 2. Correlations between inspection behavior and open field locomotion

Measure	Day of testing		
	1	2	3
Percentage of time inspecting	0.44*	0.11	0.05
Frequency of inspections	0.48**	0.12	0.05
Percentage of time spent near predator	0.43*	-0.19	0.15

Notes: * $p < 0.05$; ** $p < 0.01$.

boldness-related initial inspection and vigilance-related surveillance inspection.

We hypothesize that predator inspection would be initiated by the boldest individuals in the school. However, subsequent inspections, associated with vigilance, would not necessarily be performed by these individuals. This has important implications for between-population differences in boldness and predator inspection found in many fish species (Magurran *et al.*, 1992, 1995): fish in populations with high predation pressure have lower boldness but higher propensity to predator inspection. It is likely, that the high inspection tendency of fish from heavily predated populations is based on the motivational mechanisms linked with vigilance rather than with boldness (i.e., it may be unrelated to the propensity to take risks). The data indicating that fish from high-risk populations show signs of shyness during their predator inspection visits (e.g., they avoid the dangerous mouth region of the predator, inspect mainly in groups rather than individually, keep at a greater distance from the predator, etc., see Magurran and Seghers, 1990, 1994) seems to agree with this hypothesis. However, further experiments comparing boldness-related and surveillance predator inspection in fish from populations with different predation pressure are necessary.

ACKNOWLEDGMENTS

We thank Yu.B. Manteifel for helpful comments on an earlier draft of the paper. The first author (SB) was supported by the Russian Science Support Foundation.

This study was supported by the Russian Foundation for Basic Research (project 03-04-48789).

REFERENCES

- Altbacker, V. and Csanyi, V., The Role of Eyespots in Predator Recognition and Antipredator Behaviour of the Paradise Fish *Macropodus opercularis* L., *Ethology*, 1990, vol. 85, pp. 51–57.
- Birke, L.J.A. and Archer, J., Some Problems and Issues in the Study of Animal Exploration, in *Exploration in Animals and Humans*, Archer J. and Birke, L.J.A., Eds., New York: Van Nostrand, 1983, pp. 1–21.
- Budaev, S.V., Alternative Styles in the European Wrasse, *Symphodus ocellatus*: Boldness-Related Schooling Tendency, *Environ. Biol. Fish.*, 1997a, vol. 49, pp. 71–78.
- Budaev, S.V., “Personality” in the Guppy (*Poecilia reticulata*): A Correlational Study of Exploratory Behavior and Social Tendency, *J. Comp. Psychol.*, 1997b, vol. 111, pp. 399–411.
- Budaev, S.V., Zworykin, D.D., and Mochev, A.D., Individual Differences in Parental Care and Behavioural Profile, in the Convict Cichlid: A Correlation Study, *Anim. Behav.*, 1999, vol. 58, pp. 195–202.
- Cowan, P.E., Exploration in Small Mammals: Ethology and Ecology, in *Exploration in Animals and Humans*, Archer J. and Birke, L.J.A., Eds, New York: Van Nostrand, 1983, pp. 147–175.
- Csanyi, V., Ethological Analysis of Predator Avoidance by Paradise Fish (*Macropodus opercularis* L.). I. Recognition and Learning of Predators, *Behaviour*, 1985, vol. 92, pp. 227–240.
- Csanyi, V., Cszimada, G., and Miklosi, A., Long-Term Memory and Recognition of Another Species in the Paradise Fish, *Anim. Behav.*, 1989, vol. 37, pp. 908–911.
- Dugatkin, L.A. and Godin, J.-G.J., Prey Approaching Predators: A Cost-Benefit Perspective, *Annls Zool. Fenn.*, 1992, vol. 29, pp. 233–252.
- Godin, J.-G.J. and Davis, S., Who Dares, Benefits: Predator Approach Behaviour in the Guppy (*Poecilia reticulata*) Deters Predator Pursuit, *Proc. R. Soc. Lond., Ser. B*, 1995, vol. 259, pp. 193–200.
- Godin, J.-G. and Dugatkin, L.A., Female Mating Preference for Bold Males in the Guppy, *Poecilia reticulata*, *Proc. Nat. Acad. Sci., CASSI U.S.A.*, 1997, vol. 93, pp. 10262–10267.
- Hasson, O., Pursuit-Deterrent Signals: Communication Between Prey and Predator, *Trends Ecol. Evol.*, 1991, vol. 6, pp. 325–329.
- Hinde, R., *Animal Behaviour*, New York: McGraw-Hill, 1970.
- Huntingford, F.-A. and Coulter, R.M., Habituation of Predator Inspection in the Three-Spined Stickleback, *Gasterosteus aculeatus* L., *J. Fish. Biol.*, 1989, vol. 35, pp. 153–154.
- Lee, E.T., *Statistical Methods for Survival Data Analysis*, New York: John Wiley, 1992.
- Licht, T., Discrimination Between Hungry and Satiated Predators: The Response of Guppies (*Poecilia reticulata*) from High and Low Predation Sites, *Ethology*, 1989, vol. 82, pp. 238–243.
- Liley, N.R. and Seghers, B.H., Factors Affecting the Morphology and Behaviour of Guppies in Trinidad, in *Function and Evolution of Behavior*, Baerends, G.P., Beer, C., and Manning, A., Eds., Oxford: Clarendon Press, 1975, pp. 92–118.
- Lima, S., On the Personal Benefits of Antipredatory Vigilance, *Anim. Behav.*, 1994, vol. 48, pp. 734–736.
- Lima, S., Back to the Basics of Antipredatory Vigilance: the Group-Size Effect, *Anim. Behav.*, 1995, vol. 49, pp. 11–20.
- Magurran, A.E., The Adaptive Significance of Schooling as an Anti-Predator Defence in Fish, *Annls Zool. Fenn.*, 1990, vol. 27, pp. 51–66.
- Magurran, A.E. and Girling, S.L., Predator Recognition and Response Habituation in Schooling Minnows, *Anim. Behav.*, 1986, vol. 34, pp. 510–518.
- Magurran, A.E. and Pitcher, T.J., Provenance, Shoal Size and the Sociobiology of Predatorevasion Behaviour in Minnow Shoals, *Proc. R. Soc. Lond., Ser. B*, 1987, vol. 229, pp. 439–465.
- Magurran, A.E. and Seghers, B.H., Population Differences in Predator Recognition and Attack Cone Avoidance in the Guppy *Poecilia reticulata*, *Anim. Behav.*, 1990, vol. 40, pp. 443–452.
- Magurran, A.E. and Seghers, B.H., Predator Inspection Behaviour Covaries with Schooling Tendency Amongst Wild Guppy, *Poecilia reticulata*, Populations in Trinidad, *Behaviour*, 1994, vol. 128, pp. 121–134.
- Magurran, A.E., Seghers, B.H., Carvalho, G.R., and Shaw, P.W., Evolution of Adaptive Variation in Antipredator Behaviour, *Mar. Behav. Physiol.*, 1992, vol. 22, pp. 29–44.
- Magurran, A.E., Seghers, B.H., Shaw, P.W., and Carvalho, G.R., The Behavioral Diversity and Evolution of Guppy, *Poecilia reticulata*, Populations in Trinidad, *Adv. Study Behav.*, 1995, vol. 24, pp. 155–202.
- McLeod, P.G. and Huntingford, F.A., Social Rank and Predator Inspection in Sticklebacks, *Anim. Behav.*, 1994, vol. 47, pp. 1238–1240.
- Pitcher, T.J., Green, D.A., and Magurran, A.E., Dicing with a Death: Predator Inspection Behaviour in Minnow Shoals, *J. Fish Biol.*, 1986, vol. 28, pp. 439–448.
- Reznick, D. and Endler, J.A., The Impact of Predation on Life History Evolution in Triadadian Guppies (*Poecilia reticulata*), *Evolution*, 1982, vol. 36, pp. 160–177.
- Sokal, R.R. and Rohlf, F.J., *Biometry*, San Francisco: Freeman, 1981.
- Wilson, D.S., Glark, A.B., Coleman, K., and Dearstyne, T., Shyness and Boldness in Humans and Other Animals, *Trends Ecol. Evol.*, 1994, vol. 9, pp. 442–446.